

NF03 – Flavon exchange – a new model for neutrino oscillations



Marc Bergevin
LLNL

September 17th, 2020




New model of neutrino oscillation

Flavon Model



- Model ([2002.12269](#)) assumes that flavor change occurs at discrete external points.
 - Flavor change assumed **as a small perturbation**
 - Rotation of states is performed using a Pauli rotation trick



Proposed oscillation alternative (Flavon)

$$\begin{cases} \langle \nu_l | H_0 | \nu_m \rangle = \delta_{lm} E_{\nu_m} \\ \langle \nu_l | H_1 | \nu_m \rangle = (1 - \delta_{lm}) f_{lm} \end{cases}$$

$$U(t) = e^{-i(f_{e\mu} \sigma_x + \Delta E \sigma_z)t/\hbar} e^{-i(E_{tot} I)t/\hbar} = U_{osc} U_{E_\nu}$$

PMNS Constant amplitude. Maximum number of neutrinos that can be lost

$$P_{ee} = | \langle \nu_e(t) | \nu_e \rangle | = 1 - \sin^2(2\theta_{12}) \sin^2\left(\frac{\Delta m_{12}^2 t}{4E_\nu}\right)$$

Amplitude not constant: Energy dependence.

Flavon

$$P_{ee} = 1 - \frac{f_{e\mu}^2}{\left(\frac{\Delta m_{\nu_{\mu e}}^2}{4E_\nu}\right)^2 + f_{e\mu}^2} \sin^2\left(t \sqrt{\left(\frac{\Delta m_{\nu_{\mu e}}^2}{4E_\nu}\right)^2 + f_{e\mu}^2}\right)$$

Both formula consistent except at short and long period.
 Flavon model: Long period ($P_{ee} \sim 1 - f_{e\mu}^2 t^2$) Short period ($P_{ee} \sim 1/E_\nu^2$)

Formula untested but **consistent** with PNMS
 (except at short or long baselines)

Reactor antineutrino rate picture

Flavon vs PNMS model

Perturbation


$$f_{e\tau} = 2.49 \times 10^{-11} \text{ eV}$$

$$f_{e\mu} = 3.31 \times 10^{-14} \text{ eV}$$

$$f_{\mu\tau} = 9.65 \times 10^{-12} \text{ eV}$$

$$f_{e\tau} = 1.40 \times 10^{-10} \text{ eV}$$

$$f_{e\mu} = 2.27 \times 10^{-13} \text{ eV}$$



Provides agreement at short and long range.
Mid range experiments (AIT/JUNO) will be sensitive to this model.

New model of neutrino oscillation




Spectral agreement for Daya Bay and KamLand

Perturbation

$f_{e\tau} = 2.49 \times 10^{-11} \text{ eV}$
 $f_{e\tau} = 1.40 \times 10^{-10} \text{ eV}$

$f_{e\mu} = 3.31 \times 10^{-14} \text{ eV}$
 $f_{e\mu} = 2.27 \times 10^{-13} \text{ eV}$

$f_{\mu\tau} = 9.65 \times 10^{-12} \text{ eV}$



Reproduces expected features for flagship Daya Bay and KamLand experiments

New model of neutrino oscillations

Reproduces anomalous experimental results, but predicts CP Violation

Perturbation

$$f_{e\tau} = 2.49 \times 10^{-11} \text{ eV}$$


$$f_{e\mu} = 3.31 \times 10^{-14} \text{ eV}$$

$$f_{\mu\tau} = 9.65 \times 10^{-12} \text{ eV}$$


$$f_{e\tau} = 1.40 \times 10^{-10} \text{ eV}$$

$$f_{e\mu} = 2.27 \times 10^{-13} \text{ eV}$$


e-neutrinos and **e-antineutrinos** oscillate differently



e-neutrinos and **μ-antineutrinos** share parameters



μ-neutrinos and **e-antineutrinos** share parameters




Reproduces MSW effect and reconciles with SAGE/GALLEX.

Provides adequate agreement to LSND/MiniBoone with reactor antineutrino and solar neutrino parameters.

Reactor antineutrinos at AIT-NEO

Flavon vs PMNS model

- The Advanced Instrumentation Testbed (AIT)* is a proposed neutrino physics experimentation site at the Boulby mine in the UK. The nearest reactor is 26 km away
- The first experiment at AIT, Neutrino Experiment One (NEO) will consist of a kiloton-scale detector with a $\text{Gd-H}_2\text{O}^+$ or WbLS^\ddagger target
- The rate and spectrum difference between the PMNS and flavon models is pronounced at this standoff, regardless of fill
- **Gd-H₂O option:**
 - At 26 km standoff the flavon model predicts ~ 4 evts per month for a $\text{Gd-H}_2\text{O}$ target
 - while PMNS predicts ~ 25 evts per month
 - A $\text{Gd-H}_2\text{O}$ fill in NEO would rule out the PMNS model at 2.7σ in 1 month.
 - A $\text{Gd-H}_2\text{O}$ fill in NEO would confirm the flavon model at 5σ in 6.5 months
- **WbLS option:**
 - Not yet studied in detail, but we qualitatively expect improved energy resolution and higher rates (due to lower energy threshold)





* AIT: SNOWMASS21-NF7_NF10-IF2_IF9_Adam_Bernstein-096.pdf

† Gd-H₂O: SNOWMASS21-NF7_NF10-IF9_IF0_Adam_Bernstein-097.pdf

‡ WbLS: SNOWMASS21-NF7_NF10-IF9_IF0_Adam_Bernstein-098.pdf

Conclusion

- I'm putting forward a new model for neutrino mixing – the flavon model
- The flavon model can be tested at two upcoming antineutrino reactor experiments (JUNO and AIT-NEO)
 - The AIT-NEO program in the Boulby mine is a new, highly leveraged resource for the HEP community to pursue technology and physics
 - JUNO is predicted to observe a rate and spectrum that will also refute or confirm the flavon model.





Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.